
Effects of Cladribine Tablets on MRI Outcomes in High Disease Activity (HDA) Patients with Relapsing Multiple Sclerosis (RMS) in the CLARITY Study

G. Giovannoni¹, K. Rammohan², S. Cook³, G. Comi⁴, P. Rieckmann⁵, P. Soelberg-Sorensen⁶, P. Vermersch⁷, F. Dangond⁸, C. Hicking⁹

1 lestitute, Barts and The London, Blizard Institute of New Jersey, New Jersey San Raffaele, Ospedale San Raffaele, Milan, Italy; Department of Neurology, University of Copenhagen, Billerica, MA, USA; Medical Park Loipl and University of Copenhagen, Billerica, MA, USA; Medical Park Loipl and University of Copenhagen, Billerica, MA, USA; Medical Park Loipl and University of Copenhagen, Billerica, MA, USA; Medical Park Loipl and University of Copenhagen, Billerica, MA, USA; Medical Park Loipl and University of Copenhagen, Billerica, MA, USA; Medical Park Loipl and University of Copenhagen, Billerica, MA, USA; Medical Park Loipl and University of Copenhagen, Billerica, MA, USA; Medical Park Loipl and University of Copenhagen, Billerica, MA, USA; Managen, MA, USA; Managen, MA, USA; Managen, MA, USA; Managen, MA, USA; MA, USA; Managen, MA, USA; MA, USA;

INTRODUCTION

- CLARITY was a large (1326 patients), randomized, double-blind study in patients with early and moderately advanced active relapsing multiple sclerosis (RMS) (see Figure 1 for study design), in which treatment with cladribine tablets showed strong efficacy versus placebo (PBO) over 2 years.1
- Patients with RMS who show an increased rate of relapse or disability progression can be described as having high disease activity (HDA). Analysis of CLARITY may provide insights into the efficacy of cladribine tablets in patients with evidence of HDA.



OBJECTIVE

MRI, magnetic resonance imaging; RMS, relapsing multiple sclerosis.

 To compare the effects of cladribine tablets 3.5 mg/kg versus placebo on outcomes assessed by magnetic resonance imaging (MRI) in subgroups of CLARITY patients with evidence of HDA at study entry, using two HDA definitions.

METHODS

- The CLARITY study enrolled patients aged 18–65 years with a definite diagnosis of relapsing remitting multiple sclerosis (RRMS):2
- ≥ 1 relapse in the 12 months before study entry, but no relapses within the 28 days before entry.
- neurological lesions detectable by MRI consistent with MS.
- an expanded disability status scale (EDSS) score of 0−≤ 5.5.
- Patients were excluded if they had received a disease-modifying drug (DMD) within 3 months before study entry, or if treatment with > 1 DMD had failed.
- CLARITY patients randomized to cladribine tablets 3.5 mg/kg (N = 433) or placebo (N = 437) were retrospectively analyzed using two different HDA definitions based on relapse history, prior treatment, and MRI characteristics.

- Two overlapping sets of criteria (Figure 2) were applied in the analysis of baseline disease characteristics to subdivide patients into HDA groups based upon:
- 1. High relapse activity (HRA); patients with ≥ 2 relapses in the year before study entry, regardless of prior use of DMD.
- 2. HRA plus disease activity on treatment (HRA+DAT); patients with ≥ 2 relapses in the year before study entry, regardless of prior use of DMD, plus patients with ≥ 1 relapse AND ≥ 1 T1 gadolinium (Gd+) or ≥ 9 T2 lesions in the year before study entry while on DMD therapy.
- The ability of these criteria to retrospectively identify patients with HDA and the effects of cladribine tablets 3.5 mg/kg versus placebo on relapse rates and disability outcomes in these patients has been presented previously.^{3,4}

igure 2. Definitions of High Disease Activity HRA+DAT subgroup (n = 289) HRA subgroup (n = 261) (n = 261)(n = 261)Patients with ≥ 2 relapses in the year Patients with ≥ 2 relapses in the year before study entry, regardless of before study entry, regardless of treatment status treatment status Naïve. n = 187 Naïve. n = 187 Treated, n = 74 • Treated. n = 74 (n = 28)Patients with ≥ 1 relapse AND ≥ 1 T1 Gd+ or ≥ 9 T2 lesions in the year before study entry while on therapy with other DMDs

NB These definitions are not exclusive; to a large extent, the subgroups comprise the same patients. DAT, disease activity on treatment; DMD, disease modifying drug; Gd+, gadolinium-enhancing; HRA, high relapse activity


- Efficacy analyses from CLARITY assessed the overall population and HDA subgroups and corresponding non-HDA subgroups based on cumulative numbers of new T1 Gd+ lesions, cumulative numbers of active T2 lesions, and cumulative numbers of combined unique active (CUA) lesions (defined as a new T1 Gd+ lesion, or a new or enlarging T2 lesion, or both, without double-counting).
- All HDA analyses were post hoc and not pre-specified; no multiplicity adjustments were done to the resulting P values. All comparisons where the P value was less than 0.05 by statistical testing should be regarded as nominally significant.

RESULTS

Patients

- Demographics and disease characteristics were similar across the subgroups of patients who met the HDA definitions (**Table 1**). The demographics and disease characteristics of patients who did not fulfill the HDA criteria were also similar across the subgroups (data not shown).
- The overall analysis involved 870 patients randomized to placebo (n = 437) or cladribine tablets 3.5 mg/kg (n = 433):
- Among the non-HRA patients, 306 received placebo and 303 received cladribine tablets 3.5 mg/kg. - In those who did not meet the HRA+DAT criteria (i.e. non-HDA patients), 288 received placebo and 293 received cladribine tablets 3.5 mg/kg.

Table 1. Patient Demographics and Disease Characteristics by HDA Subgroup Cladribine Tablets 3.5 mg/kg (n = 131)36.7 (10.3) Age, years; mean (SD) 82 (62.6) 94 (63.1) Female, n (%)



30.0 (15.0; 39.0) | 29.9 (16.0; 39.0) | 25.3 (12.0; 33.0) | 25.2 (12.0; 33.5)


DAT, disease activity on treatment; **DMD**, disease modifying drug; **EDSS**, Expanded Disability Status Scale; **Gd+**, gadolinium-enhancing; HDA, high disease activity; HRA, high relapse activity; IQR, interquartile range; SD, standard deviation.

Cumulative New T1 Gd+ Lesions

Number of T2 lesions; mean (IQR)

- The relative risk of cumulative new T1 Gd+ lesions for patients in both subgroups treated with cladribine tablets 3.5 mg/kg over 2 years was low, with strong effects observed in each treatment subgroup (Figure 3).
- This was supportive of the risk reduction seen in the overall population.
- Neither the HRA or HRA+DAT subgroups showed statistical significance versus patients who had failed to meet the respective HDA criteria (i.e. non-HRA and non-HRA+DAT patients).

igure 3. Forest Plot of Relative Risk of Cumulative New T1 Gd+ Lesions by HDA Subgroup for Cladribine Tablets 3.5 mg/kg versus Placebo



CI, confidence interval; DAT, disease activity on treatment; Gd+, gadolinium-enhancing; HDA, high disease activity; HRA, high relapse activity; RR, relative risk.

Active T2 Lesions


(n = 140)

36.3 (9.5)

102 (72.9)

- The relative risk of active T2 lesions for patients in both subgroups treated with cladribine tablets 3.5 mg/kg over 2 years was low, with strong effects observed in each treatment subgroup (Figure 4).
- This was supportive of the risk reduction seen in the overall population.
- Neither the HRA or HRA+DAT subgroups showed statistical significance versus patients who had failed to meet the respective HDA criteria.

Figure 4. Forest Plot of Relative Risk of Active T2 Lesions by HDA Subgroup for Cladribine Tablets 3.5 mg/kg versus Placebo




CI, confidence interval; DAT, disease activity on treatment; HDA, high disease activity; HRA, high relapse activity; RR, relative risk.

Cumulative CUA Lesions

- The relative risk of cumulative CUA lesions for patients in both subgroups treated with cladribine tablets 3.5 mg/kg over 2 years was low, with strong effects observed in each treatment subgroup (Figure 5)
- This was supportive of the risk reduction seen in the overall population.
- Neither the HRA or HRA+DAT subgroups showed statistical significance versus patients who had failed to meet the respective HDA criteria.

Figure 5. Forest Plot of Relative Risk of Cumulative CUA Lesions by HDA Subgroup for Cladribine Tablets 3.5 mg/kg versus Placebo



CI, confidence interval; CUA, combined unique active; DAT, disease activity on treatment; HDA, high disease activity; HRA, high relapse activity; **RR**. relative risk.

CONCLUSIONS

- In patients with RMS in the CLARITY study, treatment with cladribine tablets 3.5 mg/kg produced statistically significant effects versus placebo on cumulative new T1 Gd+ lesions, active T2 lesions, and cumulative CUA lesions in two subgroups of patients selected using HDA criteria.
- In each HDA subgroup, there was a consistent treatment effect on radiological outcomes with cladribine tablets 3.5 mg/kg irrespective of disease activity before entry to the CLARITY study, or prior DMD use.

REFERENCES

- 1. Giovannoni G, et al. *N Engl J Med.* 2010;362:416-426.
- 2. McDonald W, et al. Ann Neurol. 2001;50:121-127.
- 3. Giovannoni G, et al. *Neurology*. 2017;88(16 suppl):P6.351.
- 4. Giovannoni G, et al. *Neurology*. 2017;88(16 suppl):P6.360.

ACKNOWLEDGMENTS

This study was sponsored by EMD Serono, Inc., a business of Merck KGaA, Darmstadt, Germany (in the USA), and Merck Serono SA – Geneva, an affiliate of Merck KGaA, Darmstadt, Germany (ROW). The authors would like to thank patients and their families, investigators, co-investigators, and the study teams at each of the participating centers and at Merck KGaA, Darmstadt, Germany. Medical writing assistance was provided by Jackie Campbell of inScience Communications, Springer Healthcare, Chester, UK, and was funded by Merck KGaA, Darmstadt, Germany.

DISCLOSURES

GG has received speaker honoraria and consulting fees from AbbVie, Atara Bio, Almirall, Bayer Schering Pharma, Biogen Idec FivePrime, GlaxoSmithKline, GW Pharma, Merck, Pfizer Inc, Protein Discovery Laboratories, Teva Pharmaceutical Industries Ltd, Sanofi-Genzyme, UCB, Vertex Pharmaceuticals, Ironwood, and Novartis; and has received research support unrelated to this study from Biogen Idec, Merck, Novartis, and Ironwood. KR has received honoraria for lectures and steering committee meetings from EMD Serono, Biogen Idec, Sanofi-Aventis, Genzyme, Novartis, Teva Neurosciences, Acorda, and Roche/Genentech. SC has received honoraria for lectures/consultations from Merck, Bayer HealthCare, Sanofi-Aventis, Neurology Reviews, Biogen Idec, Teva Pharmaceuticals, and Actinobac Biomed Inc.; has served on advisory boards for Bayer HealthCare, Merck, Actinobac Biomed, Teva Pharmaceuticals, and Biogen Idec; and received grant support from Bayer HealthCare. GC has received consulting fees from Novartis, Teva Pharmaceutical Industries Ltd., Sanofi-Aventis, Merck, Receptos, Biogen Idec, Genentech-Roche, and Bayer Schering; lecture fees from Novartis, Teva Pharmaceutical Ind. Ltd., Sanofi-Aventis, Merck, Biogen Dompè, Bayer Schering, and Serono Symposia International Foundation; and trial grant support from Novartis, Teva Pharmaceutical Ind. Ltd., Sanofi-Aventis, Receptos, Biogen Idec, Genentech-Roche, Merck, Biogen Dompè, and Bayer Schering. PR has received honoraria for lectures/ steering committee meetings from Merck, Biogen Idec, Bayer Schering Pharma, Boehringer-Ingelheim, Sanofi-Aventis, Genzyme, Novartis, Teva Pharmaceutical Industries, and Serono Symposia International Foundation. **PS-S** has served on advisory boards for Biogen, Merck, Novartis, Teva, MedDay Pharmaceuticals, and GSK; on steering committees or independent data monitoring boards in trials sponsored by Merck, Teva, GSK, and Novartis; has received speaker honoraria from Biogen Idec, Merck, Teva, Sanofi-Aventis, Genzyme, and Novartis. His department has received research support from Biogen, Merck, Teva, Novartis, Roche, and Genzyme. PV has received honoraria or consulting fees from Biogen, Sanofi-Genzyme, Bayer, Novartis, Merck, Celgene, Roche, and Almirall; and research support from Biogen, Sanofi-Genzyme, Bayer, and Merck. CH is an employee of Merck KGaA, Darmstadt, Germany. FD is an employee of EMD Serono, Inc., a business of Merck KGaA, Darmstadt, Germany.

The CLARITY study: NCT00213135.

Cladribine tablet(s) is an investigational product for the treatment of multiple sclerosis and has not been approved by the FDA for any indication in the U.S.

> Copies of this poster obtained through QR (Quick Response) code are for personal use only and may not be reproduced without written permission of the authors



GET POSTER PDF